



# GOLDEN STATE ORTHOPEDICS & SPINE

## REVERSE SHOULDER ARTHROPLASTY PROTOCOL

### **OVERVIEW**

- **Sling** – Simple sling for 5 weeks as follows:
  - Full time x 4 weeks: After week 4, begin to wean during day as tolerated
  - Continue to wear while sleeping until Week 5
- **When Supine:** For first 5 weeks, place small pillow/roll behind elbow to avoid hyperextension
  - Prevents tension on subscap (if repaired)
  - Instruct patient they should be able to visualize elbow

### **Other Precautions:**

- No lifting (greater than coffee mug) or strengthening for first 5-6 weeks
  - OK to use operative arm for activities of daily living (brushing teeth, eating, etc)
- No strengthening for first 5-6 weeks. Focus on PROM, followed by AAROM and AROM
- No pulleys x 6 weeks
- Reinforce weight limit of no greater than 10-15 lbs on operative shoulder/extremity

### **PHASE 1: Protected PROM/AAROM (Weeks 0-5)**

#### **Early Phase 1 (Weeks 0-4)**

- Sling full-time (except with therapy or when showering) for 4 weeks
- Patient will see therapist on post-op day #1 (in hospital) to learn:
  - **Pendulums** as tolerated
  - **Passive supine forward elevation** as tolerated
    - » Begin PROM in supine position with rotator cuff repairs for good scapular stabilization
  - **Gentle ER in scapular plane** to available PROM (no more than 30°)
- Begin Outpatient Therapy at 10-14 days post-op
- **Cryotherapy** (+ other soft tissue modalities): For swelling, pain and inflammation
- Supine forward flexion (initially PROM, progress to AAROM as tolerated)
- AROM of elbow, wrist, hand and neck
- Begin scapular isometrics (scapular sets – primarily retraction)
- Work on PROM in all planes (except limit ER to 30°, NO extension)
- NO weight-bearing on operative extremity for first 4 weeks
- NO Pulley exercises

#### **Late Phase 1 (Week 5)**

- **Sling:** Worn only when sleeping, wean during the day over Week 5
- Begin and progress from PROM to gentle **active-assisted ROM** as tolerated
  - Focus on assisted forward flexion <120°, ER <30° and Abd <45° in plane of scapula

### **PHASE 2: AAROM/AROM (Weeks 5-8)**

- Improve PROM and AAROM gradually into full ER in neutral, elevation <140°, IR as tolerated
- Discontinue sling after Week 5: Encourage natural arm swing
- Begin AROM: Initiate glenohumeral joint mobility primarily in elevation



# GOLDEN STATE ORTHOPEDICS & SPINE

## REVERSE SHOULDER ARTHROPLASTY PROTOCOL

- Begin AROM exercise against gravity in elevation, ER, IR behind back and abduction
- Avoid PROM with stretching in adduction past neutral, IR behind back & flexion > 120°
- Also avoid PROM/stretching with combined ER and abduction
- Encourage increased functional use of arm for ADLs
- **Begin pulleys** for forward elevation in plane of scapula (as long as > than 90° PROM)
- Begin sub-maximal deltoid isometrics (anterior, lateral & posterior) in neutral (at Week 8)
- Begin scapular strengthening exercises as appropriate (at Week 8)
- Begin assisted horizontal adduction
- Gentle glenohumeral & scapulothoracic joint mobilization, rhythmic stabilization

### **PHASE 3: AROM/ Strengthening (Weeks 9-13+)**

- Progress AROM exercise and continue PROM/stretch as needed.
- Continue to improve glenohumeral joint mobility in elevation and ER
- **Continue Strengthening:** Focus on **deltoid muscle balance** and **functional strength**
  - Utilize pool exercise program, low resistance T-band or light weights
  - Include teres minor and subscapularis (if intact per surgeon op report) strengthening
  - Progress from **submaximal isometrics** to **limited-range** to **full-range isotonic**
  - Resistive exercise below shoulder height encouraged
  - ER strength typically somewhat compromised due to underlying cuff dysfunction. Avoid overload of teres minor when strengthening.
- Maximize strength of shoulder girdle and upper extremity for light daily ADLs
- NO weight-lifting above shoulder height or lifting with weight > 10 lbs (unless instructed by surgeon)
- Avoid forceful AAROM in flexion > 140°, ER > 45°, IR behind body, horizontal adduction past neutral.

### **SAMPLE EXERCISES**

- 1) **Passive Supine Flexion/Forward Elevation:** Lie on your back. Grasp wrist with non-op hand and passively raise operative arm overhead. Aim to get to 90° by 3 weeks. In week 4, progress to 120°. Then full ROM after 6 weeks. Keep elbow bent and relaxed. Repeat 10 reps, 2-3 times/day





# GOLDEN STATE ORTHOPEDICS & SPINE

## REVERSE SHOULDER ARTHROPLASTY PROTOCOL

- 1) **Passive Pendulum Exercise:** Hold onto a chair back with non-op hand and bend forward. Let the operative arm hang down passively. Use body to passively swing arm: Forward, backward, side to side and in small circles Repeat throughout the day as tolerated.



© Healthwise, Incorporated